

IJBHS 2009120/5404

A faecal survey of gastrointestinal parasites of ruminants on the University of Maiduguri Research Farm

A. A. Biu^{*1}, A. Maimunatu², A. F. Salamatu² and E. T. Agbadu³

¹Faculty of Veterinary Medicine, PO Box 8136, University of Maiduguri, Maiduguri, Nigeria

²Department of Biological Sciences, Faculty of Science, University of Maiduguri, Maiduguri, Nigeria

³Department of Parasitology, PO Box 269, Vom Plateau State, Nigeria

(Received October 15, 2009)

ABSTRACT: Faecal survey of ova/oocysts of gastrointestinal parasites of ruminants on the University of Maiduguri research farm was conducted between January and November 2006 using saturated sodium chloride floatation and formol ether sedimentation techniques. A prevalence rate of 47.0%, 54.0% and 58.0% was obtained for cattle, sheep and goats respectively ($p>0.05$). The younger ruminants were more infected (cattle: 50.0%; sheep: 54.7%; and goats: 58.1%) compared to the older ruminants (cattle: 44.0%, sheep: 52.8% and goats: 57.9%) ($p>0.05$). Female ruminants were also more infected (cattle: 52.0%; sheep: 60.4%, and goats: 62.7%) compared to the males (cattle: 52.0%; sheep: 46.8% and goats: 51.2%) ($p>0.05$). Among cattle breeds, the Wadara had 46.9% infection while the Simmental had 50.0% ($p>0.05$). Among sheep breeds the Yankasa, Ouda and Sudanese had 53.8%, 43.8% and 100.0% infection rates respectively ($p>0.05$). Goat breeds of Borno white and Sokoto red had 54.2% and 59.6% infection rates respectively ($p>0.05$).

Keywords: Faecal, Survey, Gastrointestinal parasites, Ruminants.

Introduction

Parasitic gastroenteritis has been noted as major constraint to ruminants' productivity in terms of pathology and economic importance (Biu, *et al.*, 2006) on a clinical and subclinical level (Martinez-Gonzalez *et al.*, 1998). Although losses due to parasitic gastro enteritis have been estimated and reported in some parts of Nigeria (Akerejola *et al.*, 1979), very few studies had been conducted in Borno State (Ahmed *et al.*, 1993), which is a major livestock producing area with an estimated ruminant population put at 40% of the 70,590,000 national count (Bourn *et al.* 1994). This study was motivated by the need to isolate and identify the gastrointestinal parasites of sheep goats and cattle of the research farm in the objective of determining their prevalence.

*To whom correspondence should be addressed.
E-mail: biuvet@yahoo.com

Materials and Methods

Description of study site: The University of Maiduguri research farm lies between the Sudan and Sahel vegetation zones characterized by a short rainy season from June to September, a prolonged dry period between November and May (Udoh, 1981).

Sample collection: Faecal samples were obtained from the rectum of a hundred (100) each of randomly sampled cattle, sheep and goats kept under the traditional husbandry system of semi intensive management of the University Research farm.

Faecal examination: Faecal samples obtained were examined using saturated sodium chloride floatation and formol ether sedimentation techniques and eggs or oocysts identified using the light microscope at x40 objective and the eggs or oocysts per gram of faeces determined using the modified Mc Master technique (Hansen and Perry, 1990).

Statistical analysis: Data obtained based on age, sex and breed of the ruminants examined were analyzed using the students "t" test with p values equal to or less than 0.05 regarded as significant (Graph pad Instat, 2000).

Results

The prevalence of parasitic gastro intestinal ova /oocysts identified on the research farm as shown in Table 1 indicated an overall rate of 51.3% representing 154 infested cases out of the 300 ruminants examined. Single infection by *Eimeria* oocysts and Strongyle ova had the highest prevalence of 53 (34.4%) and 48 (31.2%) respectively, followed by *Strongyloides* ova with 17 (11.0%) while *Taenia* and Trichuriid ova had the lowest with 4 (2.6%) each ($p < 0.05$).

Mixed infections by *Eimeria* oocysts + Strongyle ova had the highest prevalence of 13 (8.4%), while the lowest was by Strongyle + *Eimeria* + *Strongyloides* and *Taenia* + *Eimeria* with 2(1.3%) each ($p < 0.05$).

Table 2 shows the prevalence of gastro-intestinal parasites of cattle examined based on breed sex, and age. Of the 100 cattle examined 47 (47.0%) were infected. No significant difference ($p > 0.05$) occurred amongst sex, age and breeds.

Table 3 shows the prevalence of gastro-intestinal parasites of sheep examined based on breed sex, and age. No significant difference ($p > 0.05$) occurred amongst sex, age and breeds.

Table 4 shows the prevalence of gastro intestinal parasites of goats examined based on breed, sex and age. An overall prevalence of 58.0% was observed with no significant difference among age, breed and sex ($p > 0.05$).

Discussion

The results of this study show that ruminants of the University of Maiduguri research farm are commonly infested with a variety of gastro intestinal parasite species with a high prevalence of 51.3% and generally very high egg/oocyst count. This finding agrees with the reports by Adejinmi and Harrison, (1997) that the incidence of parasitic gastro-enteritis of ruminants is usually high especially those kept under traditional methods of husbandry, with insidious effects that undermine host health particularly when compounded by additional stress such as malnutrition, (Pal and Qayyum, 1993), a common penalty in this semi arid region of northeastern Nigeria (Biu *et al*, 2006).

This study also revealed *Eimeria* and strongyle species as the most common parasites of the research farm ruminants, and are reported as the most incriminated gastro- intestinal parasites of domestic ruminants (Eysker and Ogunsusi, 1980) especially from other parts of Maiduguri Municipal (Biu and Eteng 2001).

Table 1: Prevalence of faecal ova/oocysts of ruminants on the University of Maiduguri research farm

Parasite ova/oocysts	No (%) of animals infected. (N = 300)
Overall	154 (51.3)
Strongyle	48 (31.2)
<i>Strongyloides</i>	17 (1.0)
<i>Trichuris</i>	4 (2.6)
<i>Eimeria</i>	53 (34.4)
<i>Taenia</i>	4 (2.6)
Strongyle + <i>Strongyloides</i>	4 (2.6)
<i>Eimeria</i> + Strongyle	13 (8.4)
Strongyle + <i>Eimeria</i> + <i>Strongyloides</i>	2 (1.3)
<i>Taenia</i> + <i>Eimeria</i>	2 (1.3)
<i>Taenia</i> + Strongyle	3 (1.9)
<i>Strongyloides</i> + <i>Eimeria</i>	4 (2.6)

Table 2: Prevalence of gastro-intestinal parasites of cattle examined based on breed, sex and age.

	No. of cattle examined	No (%) infested	Mean \pm SD (EPG)
Overall	100	47 (47.0)	6200 \pm 8626.7
Sex			
Male	50	21 (42.0)	5100 \pm 6929.6
Female	50	26 (52.0)	6200 \pm 8626.7
Age (Years)			
Younger (1 – 3.5)	46	23 (50.0)	5950 \pm 7566.0
Older (4 – 12)	54	24 (44.4)	5500 \pm 7071.1
Breed			
Wadara	96	45 (46.9)	6250 \pm 8556.0
Simmental	04	02 (50.0)	3500 \pm 4808.3

Table3: Prevalence of gastro intestinal parasites of sheep examined based on breed, sex and age.

	No. of sheep examined	No (%) infested	Mean \pm SD (EPG)
Overall	100	54 (54.0)	6500 \pm 8909.5
Sex			
Male	47	22 (46.8)	5700 \pm 7636.8
Female	53	32 (60.4)	6500 \pm 8909.5
Age (Years)			
Younger (1 – 2.5)	64	35 (54.7)	6500 \pm 8909.5
Older (3 – 4)	36	19 (52.8)	2150 \pm 2757.7
Breed			
Yankasa	80	43 (53.8)	6500 \pm 8909.5
Uda	16	07 (43.8)	3950 \pm 5161.9
Sudanese	04	04 (100.0)	7000 \pm 5798.3

Table 4: Prevalence of gastro intestinal parasites of goats examined based on the breed, sex and age.

	No. of goats examined	No (%) infested	Mean \pm SD (EPG)
Overall	100	58 (58.0)	6650 \pm 9263.1
Sex			
Male	41	21 (51.2)	5550 \pm 7566.0
Female	59	37 (62.7)	6650 \pm 9263.1
Age (Years)			
Younger (1 – 2.5)	62	35 (56.5)	6650 \pm 9263.1
Older (3 – 4.0)	38	22 (57.9)	5500 \pm 5586.1
Breed			
Borno White	48	26 (54.2)	6700 \pm 9192.4
Sokoto Red	52	31 (59.6)	4100 \pm 5656.9

Though, this study has not revealed any significant difference in infestation and egg output among age, sex and breed of the ruminants investigated, it has been reported, (Pal and Qayyum 1993), that prevalence of parasitic gastro-enteritis is related to the agro- climatic conditions like quantity and quality of pasture, temperature, humidity and grazing behaviour of the host and that susceptibility to infestation is influence by factor of age (Richard *et al* 1990); breed (Prolamkarn *et al* 1997);species (Vlassoff *et al* 1997); health status and physiological factors of pregnancy and previous exposure to parasites (Bekele *et al* 1987).

In conclusion, the egg output by gastrointestinal parasites; which is an important index to determine the degree of pasture contamination with parasitic ova/oocysts suggest periods of strategic treatment for the

research farm ruminants and although effective chemotherapy is available for most gastrointestinal parasites, because of their complex epizootiology it is paradoxically simple to treat the individual but almost impossible to control the disease (Allonby 1980) under the semi intensive management system of the research farm.

References

- Adejinmi, J.O. and Harrison, L.J.S. (1997). Parasitic nematodes of domestic ruminants in Nigeria: impact on ruminant production and control. *Tropical Veterinarian* 15:137-148.
- Allonby, E.W. (1980). Strategy and costs of helminth control with indications for research. In: impact of animal disease research and control on livestock production in Africa. Proceedings of the conference held in Nairobi; Sept. 1980 OAU/ILRAD ILCA/DSE, Berlin pp.135-140.
- Bekele, T., Mukasa-Mugerwa, E. and Scholtens, R.G. (1987). Seasonal changes in nematode faecal egg counts of sheep in Ethiopia. *ILCA Bulletin* 29:9-11.
- Biu, A.A. and Eteng, F.O. (2001). Some parasites causing diarrhea amongst kid goats in Maiduguri, Nigeria. In: Proceedings of the 6th Annual Conference of Animal Science Association of Nigeria. September 17th -19th, 2001 University of Maiduguri Convocation Square. Pp. 232-233.
- Bourn, D., Wint, W., Blench, R. and Woolley, E. (1994). Nigerian Livestock Resources Survey. *World Animal Review*. 78(1): 49-59.
- Cheijna, S.N. (1986). The epizootiology and control of parasitic gastro-enteritis of domestic ruminants in Nigeria. *Helminthological Abstracts*. 55:413-429.
- Eysker, M. and Ogunsusi, R. A. (1980). Observations on epidemiological and clinical aspects of gastro intestinal helminthiasis of sheep in northern Nigeria during the rainy season. *Revue Veterinary Science* 28: 58-62.
- Graphpad Software Instat version 3.05 (2000). Guide to choosing and interpreting statistical tests. Graphpad software Inc. San Diego, California USA www.com.graphpad.
- Martinez- Gonzalez, B., Diez-Banos, N. and Rojo –Vazquez, F. A. (1998). An epidemiological study of gastro intestinal parasitism in dairy sheep flocks in Leon (NW) Spain. *Small Ruminant Research*, 27: 25-30.
- Pal, R.A. and Qayyum, M. (1993). Prevalence of gastro intestinal nematodes of sheep and goats in upper Punjab Pakistan. *Pakistan Veterinary Journal* 13 (3): 138-141
- Prolamkarn, W., Pandey, V.S., Ngampongsai, W., Choldumrongkul, S., Saithanoo, S., Rattanachon, L. and Verthust, A. (1977). Genetic resistance of three genotypes of goats to experimental infection with *Haemonchus contortus*. *Veterinary Parasitology* 8: 79-90.
- Richard, S., Cabaret, J. and Cabourg, C. (1990). Genetic and environmental factors associated with nematode infection in dairy goats in Northwestern France. *Veterinary Parasitology* 36: 237-243.
- Udoh, R.K. (1981). *Geographical regions of Nigeria*. Heinmann Educational Books Ltd. Ibadan, Nigeria. 304 pp.
- Vlassoff, A., Bisset, S. A. and Mc Murtry, L.W. (1999). Faecal egg counts in Angora goats following natural or experimental challenge with nematode parasites within flock variability and repeatabilities. *Veterinary Parasitology* 84: 113-123.